

# National Agricultural Library

REG-10160241

Relais

SAM TESTA  
ARS  
MSA/NATIONAL SEDIMENTATION LABORATORY  
PO BOX 1157, 598 MCELROY DRIVE  
OXFORD, MS 38655

ATTN: SUBMITTED:  
PHONE (662) 232-2933 PRINTED: 2003-09-25 11:18:48  
:  
FAX: REQUEST NOREG-10160241  
E-MAIL SENT VIA: Manual

-----  
REG Regular Copy Journal  
-----

DELIVERY: E-mail: stesta@ars.usda.gov  
REPLY: Mail:

THIS IS NOT A BILL.

NOTICE: THIS MATERIAL MAY BE PROTECTED BY COPYRIGHT LAW

----National-Agricultural-Library/-Document-Deliver-----

E160241

QH 96

AIIS8  
-----

9/24/05 SEP23NAL05 Date Not Needed After: 12/31/03

SAM TESTA  
ARS, USDA, MSA, NATIONAL SEDIMENTATION LABORATORY  
PO BOX 1157, 598 MCELROY DRIVE  
OXFORD, MS 38655  
Patron ID: 101061

Cooper, C. M., Testa, III, S. and Shields, Jr., F. D.  
Stream Restoration: Response of Benthos to Engineered Stable Riffle/Pool Habitat. ✓  
Verh. Int. Ver. Theor. Limnol. (SIL)  
Vol. 27, Part 3, 1520-1527. 2001. 52

I have read the warning on copyright restrictions and accept full responsibility for compliance.

Maximum Cost: n/a

SAM TESTA 9/23/03 Phone# (662) 232-2933  
ARIEL IP Address: 130.74.184.144  
stesta@ars.usda.gov

-----

SEP 24 2005

## Stream restoration: response of benthos to engineered stable riffle/pool habitat

Charles M. Cooper, Sam Testa, III and F. D. Shields, Jr.

### Introduction

Habitat degradation is a worldwide phenomenon. In aquatic ecosystems channel incision is a common destructive mode of stream erosion in smaller watersheds which have little or no geologic control to maintain streambed base level. In the United States, resulting channel width increases have been reported as high as 100–1,000%, with associated sediment yields of over 1000 metric tons  $\text{km}^{-2} \text{year}^{-1}$ . Problems commonly associated with incision include elevated sediment loads, altered hydrologic characteristics resulting from isolation from the floodplain, and loss of normal instream habitat structures such as pools, riffles, woody debris, and undercut banks. Physical characteristics following bank failure and resultant channel widening include shifting bed substrates covered by very shallow and fairly uniform water depths, lack of woody riparian vegetation, and sharper hydrographs during storm events. These changes to the stream ecosystem impact on the biotic community, resulting in altered, and sometimes depauperate populations.

If unassisted, geological time scale is probably needed for recovery from incision-related changes because of the landscape scale impact. Rehabilitation measures are common in degraded stream systems, but restoration of habitats in incised warm-water sand bed streams has received little attention. In this study, we examined the changes in invertebrate community structure elicited by implementation of stabilization structures designed to provide a low-cost approach to restore incised stream channels.

### Study sites

We selected 1-km reaches in three northwest Mississippi streams, two of which were located in an agricultural setting and had experienced widespread incision (Goodwin Creek and Bobo Bayou) and one reference site which had not (Toby Tubby Creek). Toby Tubby Creek was 7–9 m wide and about 2 m deep, bordered by a forested wetland, with a total watershed area of about 38  $\text{km}^2$  (12% urban, 15%

cropland, 22% pasture, and 48% forest). Bed material was sand with  $D_{50}$  ranging from 0.06 to 0.55 mm. The stream flowed in a sinuous channel with almost total canopy, and large woody debris was common, with occasional beaver dams. Stream bed material was predominately sand, but clay and woody debris were common. Typical riffle substrates (gravel, cobble or rock) were absent, but woody debris and shallow runs interrupted the dominant pool/deep run characteristics of the stream reach. Average water depth was 45 cm. Base flow measurements averaged  $0.4 \text{ m}^3 \text{ s}^{-1}$ .

Goodwin Creek, selected as the test site, has a watershed area of approximately 21  $\text{km}^2$  (13% row crops, 60% pasture and idle lands, and 28% forest). This straightened and historically incised channel was 20–70 m wide and 4–5 m deep. The stream bed was a mixture of sand and gravel ( $D_{50}$ , 0.24–8.5 mm), but riffles were unstable and transient. At base flow, the pool habitat (depth >30 cm and velocity <10  $\text{cm s}^{-1}$ ) composed only 5–20% of the water area. The mean water surface width was approximately 6 m, and the mean depth was 20 cm. Instantaneous discharge measurements averaged  $0.06 \text{ m}^3 \text{ s}^{-1}$ . Stone riprap groins and longitudinal toe protection were placed at locations within and just upstream of the reach during 1989–1991.

Bobo Bayou, our straightened incised reference stream, has a watershed area of about 16  $\text{km}^2$  (13% row crops, 54% idle land and pasture, 33% forest). It was 25–29 m wide at the channel top, and 4.5 m deep. Stream bed sediments were sand and gravel similar to those at Goodwin Creek ( $D_{50}$ , 0.42–1.5 mm). This creek had considerable canopy cover, but woody debris in the channel was not abundant. Mean depths ranged from 5 to 17 cm at base flow, and discharges at base flow averaged  $0.01 \text{ m}^3 \text{ s}^{-1}$ .

### Rehabilitation

Rehabilitation efforts in Goodwin Creek were designed to accelerate the natural evolution of the physical attributes of the stream corridor toward

conditions typical of (SHIELDS et al. 1992), to extend existing groins to create 18 small weirs times the average base km study reach. Weir facing upstream to for channel, with the intaining scour pools for the stream reach. The 2 m wide and 0.6 m ambed except for a 1 with the stream bed upon an existing ston inserted into the low- 3 m. Stones ranged 50–85% weighing less native black willow ( long by 8–30 cm dia low-flow bank immed weir. All rehabilitation the winter of 1992–19

### Methods

Invertebrates were co regions within the 1 stream bed sediment organic matter) grab from large woody de taken at each creek du not available. Bed sed cm (2-inch) diameter cm. Three composite channel formed each were taken from sand sample consisted of ty ume of approximately twig or other vegeta (riprap) and large wo by compositing brush and collecting dislod with a standard D-f stream. No gravel or the non-incised ref Creek). Samples were rose bengal dye was samples were first el remove invertebrates only hand picked. Im fied to genus (except then enumerated. Ch to genus, but were tr the large number of it ration requirements for percent contribu

## red stable riffle/

s, Jr.

and 48% forest). Bed material ranging from 0.06 to 0.55 in a sinuous channel with large woody debris was beaver dams. Stream bed was sand, but clay and silt. Typical riffle substrates were absent, but woody debris interrupted the dominant riffle substrates of the stream reach. Mean depth was 0.45 m. Base flow measure-

ment as the test site, has a total area of approximately 21 km<sup>2</sup> (13% cropland, 28% forest). Historically incised channel was 1.5 m deep. The stream bed was sand and gravel (D<sub>50</sub>, 0.24–8.5 mm) and silt (depth >30 cm and depth <30 cm) composed only 5–20% of the stream bed. Mean water surface width was 1.5 m. Mean depth was 20 cm. Mean velocity measurements averaged 0.1 m s<sup>-1</sup>. Cross sections and longitudinal toe locations within and just upstream of the incised reference reach were measured during 1989–1991.

Goodwin Creek were natural evolution of the stream corridor toward

conditions typical of the non-incised reference (SHIELDS et al. 1992, 1994). Stone riprap was added to extend existing groins across the stream channel to create 18 small weirs at intervals approximately six times the average base flow channel width in the 1-km study reach. Weirs were v-shaped, with vertices facing upstream to focus flow into the center of the channel, with the intention of creating and maintaining scour pools followed by riffle areas through the stream reach. The crests of the weirs were about 2 m wide and 0.6 m higher than the existing streambed except for a 1-m wide central gap left level with the stream bed. One side of each weir was built upon an existing stone groin while the other was inserted into the low-flow channel bank a distance of 3 m. Stones ranged from 0.2 to 450 kg, with 50–85% weighing less than 36 kg. Ninety dormant native black willow (*Salix nigra*) posts, each 1.5 m long by 8–30 cm diameter, were planted along the low-flow bank immediately downstream from each weir. All rehabilitation efforts were completed during the winter of 1992–1993.

### Methods

Invertebrates were collected from four 50-m-long regions within the 1-km reaches. Four samples of stream bed sediments, CPOM (coarse particulate organic matter) grab samples, and surface brushings from large woody debris and introduced rock were taken at each creek during each collecting visit unless not available. Bed sediments were sampled using a 5-cm (2-inch) diameter lexan corer to a depth of 10 cm. Three composited cores taken across the stream channel formed each sample. Separate core samples were taken from sand and gravel areas. Each CPOM sample consisted of two composited grabs (total volume of approximately 700 cc) of leaf and/or small twig or other vegetative matter. Introduced rock (riprap) and large woody debris snags were sampled by compositing brushings from two 100-cm<sup>2</sup> areas and collecting dislodged material and invertebrates within a standard D-frame aquatic net held downstream. No gravel or introduced rock was present at the non-incised reference stream (Toby Tubby Creek). Samples were preserved in 80% ethanol and rose bengal dye was added to aid in sorting. Core samples were first elutriated, then hand picked to remove invertebrates. All other sample types were only hand picked. Invertebrates were usually identified to genus (except Chironomidae: Diptera) and then enumerated. Chironomidae were not identified to genus, but were treated as a single taxon due to the large number of individuals and time and preparation requirements for determination. Thus, values for percent contribution of the dominant taxon

(PCDT) do not include counts of Chironomidae. Data from the four sampling reaches within each creek were composited by sample type and date for each creek to reduce the effects of intra-sample type variation (HARRIS et al. 1995). Subsequent observations are reported from yearly averages computed from both spring and fall sample information unless otherwise indicated to remove seasonal bias. Voucher specimens of each invertebrate taxon are retained in the laboratory reference collection.

Physical habitat characteristics were measured during benthos collection semi-annually (spring and autumn) for 2 years before (1991–1992) and during the 3 years following construction (1993–1995) at 1-km reaches in each stream. The non-incised reference stream (Toby Tubby Creek) was not sampled during spring of 1991. Physical data collection consisted of 75–140 measurements taken along 15–24 transects at each site. Measurements included depth, velocity at 0.6 times the water column depth, bed type, notes on presence and number of beaver dams or man-made structures, and area of large woody debris formations. From these data, summary statistics for pre- and post-restoration were calculated. Scour holes below each weir were measured during summer base flow using a tape and wading rod. The thalweg profile was surveyed 3 months after restoration construction activities, and compared to a profile taken during 1985.

### Results

#### Physical

Scour holes formed rapidly downstream of each weir following construction; the average dimensions were 11 m wide by 14 m long by 130 cm deep. Gravel deposits formed riffle-like features upstream of many weirs. Thalweg profiles showed a stair-step formation along the 18 weirs and associated scour holes similar to riffle-pool progressions in well-developed natural streams. Prior to the construction only four pool areas at irregular intervals were recorded, and observed fill and scour sequences dominated the habitat conditions as sand migrated downstream or was flushed by large events with excessive energy. Post-construction measurements showed local increases in water width of over 100% and local increases in depth of 0–1,000%.

In Goodwin Creek restoration activities produced changes in mean width, depth, and velocity of +56%, +148%, and -67%, respec-

tively Pool areas (>30 cm depth) increased from 20% of the water area before construction to 53% only 3 months after the restoration activities had begun. After 7 months, beaver activity to seven of the weirs increased the pool area to 75%, the water width was nearly twice the pre-construction values, and the mean water depth approached that of the non-incised reference stream, Toby Tubby Creek. The mean velocity, meanwhile, decreased sharply. High mortality of willow post plantings (-80%) was observed due to competition by Kudzu (*Pueraria lobata*) and other herbaceous species, impermeable or droughty soils, and beaver (*Castor canadensis*) activity. In spite of beaver activity, the large woody debris density in Goodwin Creek remained about an order of magnitude less than the incised reference (Bobo Bayou) and the non-incised reference stream (Toby Tubby).

Changes in the stream bed composition indicate substrate dynamics in incising channels. Goodwin Creek had a 10% decrease in the measured occurrence of sand, with an attendant increase in rock and gravel bed types and a mean particle size diameter ( $D_{50}$ ) increase from 0.55 to 0.75 mm. The incised comparison stream, Bobo Bayou, incurred larger changes in bed composition as sand substrate decreased more than 35% and large increases of gravel and debris were observed from ongoing channel incision and bank failure. The mean particle size ( $D_{50}$ ) at Bobo Bayou increased from 0.52 to 2.0 mm. Toby Tubby Creek showed little change during the study, with a small decrease in debris density exposing more clay substrate, and  $D_{50}$  (0.4 mm) remained relatively unchanged. A more comprehensive discussion of the physical habitat changes resulting from the rehabilitation efforts is given by SHIELDS et al. (1995).

*Invertebrates*

During the 5-year study 67,029 invertebrates were collected. The number of taxa (at the genus level) exhibited an increasing trend at all creeks through the first 3 years (1991-1993) of the study (possibly due to long-term recovery following extremely high flows in early 1991). During 1994 (the second year following reha-

bilitation at Goodwin), the number of taxa for all creeks was similar, but during 1995, a distinct and similar decline occurred in both the incised reference stream and the non-incised reference, while taxa at the rehabilitated stream increased. Pre- and post-rehabilitation averages show that the number of taxa increased by eight at both the rehabilitated and non-incised reference stream (-70%), but only by four (32%) at the incised reference (Table 1).

Following rehabilitation, the average number of individuals at Goodwin Creek decreased by 18% from the pre-modification value. This change resulted from a 28% decrease in the average number of Chironomidae present, while there was a 118% increase of the (smaller) non-chironomid component of invertebrates. Numbers of total individuals at the incised reference stream (Bobo Bayou) increased by 166%, and numbers at the non-incised reference (Toby Tubby) increased by 60%. Pre-modification values for the ratio of Chironomidae individuals to other individuals were: Goodwin Creek (15.3:1), Toby Tubby (5.6:1), and Bobo Bayou (6.2:1). Post-rehabilitation values were Goodwin Creek (4.6:1), Toby Tubby (3:1), and Bobo Bayou (11.8:1).

The non-chironomid portion of the fauna at Goodwin Creek before rehabilitation was dominated by three families: Hydropsychidae (Trichoptera - 36%), Simuliidae (Diptera - 28%), and Baetidae (Ephemeroptera - 20%). Other taxa present but contributing less than 10% each included Siphonuridae (8%) and Caenidae (4%) (Ephemeroptera), Annelida (3%), and Physidae snails (1%). Following rehabilitation, Simuliidae made up 50% of all individuals collected, with the next most abundant group being the Annelida (18%). Baetid mayflies decreased to 7%, and Hydropsychid caddisflies decreased to only 6%. Copepoda represented 6% of the fauna, Ceratopogonidae (Diptera) 5%, and Lymnaeid snails 3%. Caenid mayflies contributed only 1%, as did the Hydrophilidae (Coleoptera). Asselid crustacea, physid snails, coenagrionid odonates, and gammarid crustacea each contributed less than 1% to the fauna.

The number of Ephemeroptera, Plecoptera,

Table 1. Average pre- and post-rehabilitation sample data. Goodwin Creek (non-incised reference) and Toby Tubby Creek (incised reference) are compared with the average of other displayed values.

Number of taxa	
Number of individuals	
Number of EPT	
Number of Chironomidae	
Number of non-Chironomidae	
Percent contribution of Chironomidae	

and Trichoptera (EPT) at Goodwin Creek followed a similar trend. Taxa increased at both Toby Tubby (56% and 28%, respectively) and Goodwin Creek (number of taxa in the Goodwin Creek decreased 10% before rehabilitation). EPT richness increased at Goodwin Creek (from 5.5 to 7.6) and decreased at the incised reference (from 5.5 to 6.0). Percent contribution of Chironomidae (PCDT) decreased by 28% following rehabilitation at the incised stream, Toby Tubby Creek. PCDT at the incised reference stream decreased 6%.

Collector-filterer families increased dramatically following rehabilitation at Goodwin Creek. Modification yearly to

Table 1. Average pre- and post-rehabilitation values, and amount and percent change for selected invertebrate sample data. Goodwin Creek (rehabilitated stream), Bobo Bayou (incised reference), Toby Tubby Creek (non-incised reference). (Apparent incongruity of some "Change" column values is due to numerical rounding of other displayed values).

		Before	After	Change	% Change
Number of taxa	Goodwin Creek	13	21	8	65%
	Bobo Bayou	13	18	4	32%
	Toby Tubby Creek	11	19	8	78%
Number of individuals	Goodwin Creek	3367	2763	-604	-18%
	Bobo Bayou	1212	3224	2012	166%
	Toby Tubby Creek	1015	1624	609	60%
Number of EPT	Goodwin Creek	145	73	-73	-50%
	Bobo Bayou	126	160	35	28%
	Toby Tubby Creek	144	225	81	56%
Number of Chironomidae	Goodwin Creek	3142	2271	-870	-28%
	Bobo Bayou	1044	2972	1928	185%
	Toby Tubby Creek	819	1221	402	49%
Number of non-Chironomidae	Goodwin Creek	226	491	266	118%
	Bobo Bayou	168	252	84	50%
	Toby Tubby Creek	147	403	256	174%
Percent contribution dominant taxon	Goodwin Creek	63	44	-19	-30%
	Bobo Bayou	47	45	-3	-6%
	Toby Tubby Creek	49	37	-11	-23%

and Trichoptera (EPT) decreased 50% at Goodwin Creek following rehabilitation, but increased at both Toby Tubby and Bobo Bayou (56% and 28%, respectively). EPT richness (number of taxa in the EPT orders) in Goodwin Creek decreased from an average of seven taxa before rehabilitation to 6.3 afterwards. EPT richness increased at the non-incised reference (from 5.5 to 7.6, average) and incised reference (from 5.5 to 6.3, average) streams. The percent contribution of the dominant taxon (PCDT) decreased by 30% at Goodwin Creek following rehabilitation. PCDT of the reference stream, Toby Tubby, decreased 23%, and PCDT at the incised reference, Bobo Bayou, decreased 6%.

Collector-filterer functional group individuals increased dramatically during the year following rehabilitation at Goodwin, from pre-modification yearly totals of 283 and 283, to

1,423 individuals during the calendar year, 1993. The number of collector-filterer individuals for the next year was only 362, and for the final year of the study was only 13. A similar pattern was observed for the non-incised reference stream, with the total number of collector-filterer individuals expanding from 62 and 427 during the first 2 years of study, to 1,540 individuals during 1993, then decreasing to only 216 and 119 for the final 2 years of the study. The incised reference stream showed no increase during 1993 but rather a decrease in the number of collector-filterers, with observed numbers of 36, 326, 179, 277 and 6 individuals during the consecutive years of the study.

The shredder functional group exhibited a similar pattern at all three creeks, with extremely few individuals collected during the first 2 years of the study, a peak amount captured during 1993, and moderate numbers col-

n), the number of taxa for  
r, but during 1995, a dis-  
cline occurred in both the  
eam and the non-incised  
at the rehabilitated stream  
post-rehabilitation averages  
r of taxa increased by eight  
ted and non-incised refer-  
but only by four (32%) at  
(Table 1)

ation, the average number  
odwin Creek decreased by  
-modification value. This  
n a 28% decrease in the  
Chironomidae present,  
% increase of the (smaller)  
component of invertebrates.  
individuals at the incised ref-  
Bobo Bayou) increased by  
s at the non-incised refer-  
increased by 60%. Pre-  
for the ratio of Chironomi-  
other individuals were:  
3.3:1), Toby Tubby (5.6:1),  
6.2:1). Post-rehabilitation  
win Creek (4.6:1), Toby  
Bobo Bayou (11.8:1).

mid portion of the fauna at  
ore rehabilitation was domi-  
nated by: Hydropsychidae (Tri-  
mulidae (Diptera - 28%),  
phemeroptera - 20%) Other  
contributing less than 10%  
Conuridae (8%) and Cae-  
teroptera), Annelida (3%),  
1%). Following rehabilita-  
up 50% of all individu-  
the next most abundant  
nelida (18%). Baetid may-  
%, and Hydropsychid cad-  
o only 6%. Copepoda  
e fauna, Ceratopogonidae  
mnacoid snails 3%. Caenid  
d only 1%, as did the  
optera). Asselid crustacea,  
rionid odonates, and gam-  
contributed less than 1%

phemeroptera, Plecoptera,

lected during the final 2 years of the study. Peak numbers during 1993 were highest at the rehabilitated stream, Goodwin Creek, where 268 shredders were collected, while only 115 and 91 were collected at the non-incised and incised reference stream, respectively. The percent shredder individuals of the total individuals collected was lowest at the two incised streams before rehabilitation efforts, and highest at the non-incised reference site. Following rehabilitation, Goodwin Creek increased to a ratio near that of the non-incised reference in the first year, then exceeded the non-incised reference during the final 2 years.

Observed changes in the proportions of functional feeding groups collected from Goodwin Creek before and after rehabilitation are summarized in Fig. 1. Filtering collector, scraper, parasite and piercer group contributions were relatively unchanged. The gathering collector contribution decreased by 70%, while the predator contribution increased nearly 5-fold, and the shredder contribution increased by a factor of 6.

**Discussion**

The physical response to rehabilitation efforts was generally as expected. Such rapid response patterns of erosion and deposition were typical of those observed by other authors for weirs in

steeper streams, larger rivers, and in physical models. Beaver activity compounded the stream response to the weirs, creating a greater pool habitat at the expense of the riffle area. Though unplanned, this occurrence may represent a return to more natural conditions that existed prior to European settlement in the area. Beaver activity did not affect drainage or cause flooding of adjacent fields due to the water-carrying capacity of the incised creek. As noted above, beaver impoundments were common at the non-incised reference stream.

Biological interactions, especially competition, may dominate community formation and dynamics in constant environments. Physical disturbance, conversely, prevents the continuing domination by a small group of specialized competitors and allows rapid colonizers to co-exist (YODZIS 1986). Stream systems are subject to multiple disturbances that fluctuate widely in intensity and time of occurrence, including seasonal and storm-related hydraulic events and allochthonous inputs of materials from the watershed and upstream channel. Patch dynamics (PATRICK 1975) exemplifies the rapid colonization/reproduction that characterizes stream organisms evolved to inhabit these habitats. In Goodwin Creek, we observed continuing changes in the invertebrate community despite the creation of a more stable riffle-pool

sequence. We specifically note shifts to (1) erosion and (2) creation of the pool habitat with greatly enlarged habitat area. This shift in habitat structure from fish, and the ability of aquatic invertebrates to exploit this habitat.

The first half of the study was a response to a large-scale catastrophic high flow event. MITCHELL (1991) found that such events have potentially major effects on stream structure and communities in stream systems. The similarity in stream structure during the first 4 years of the study indicates that a recovery of stream structure occurred in the first 3 years and that the stream recovered by the fourth year. The rehabilitated stream structure (which decreased in pool habitat) during the first 4 years could be attributed to the increased stream width (which occurred during) some regional stream restoration projects. Species richness at the rehabilitated stream was similar to that of the non-incised reference stream.

In addition to the physical changes in stream flows, rehabilitation of stream structure (pool habitat in Goodwin Creek) resulted in a 70% increase in water velocity and a 70% decrease in sand and silt. Invertebrate abundance increased in the pool habitat change, the stream recovered rapidly. The fish biomass was 6.8 kg/100 m<sup>2</sup> in the pool habitat and 10.0 kg/100 m<sup>2</sup> in the riffle habitat. The fish biomass increased from 3.30 kg/100 m<sup>2</sup> in the pool habitat to 100 m<sup>2</sup>. Invertebrate abundance decreased rock and natural substrate. The major diet component in the diet content analysis on the fish was sunfish, *Lepomis microlophus*, and yellow bullheads (*Ictalurus nebulosus*) showed that the diet was of substrate oriented. Occasional incidents of Chironomidae were found in the diet content in all three




Fig. 1. Percent functional feeding groups before and after rehabilitation in Goodwin Creek, Mississippi

er rivers, and in physical  
y compounded the stream  
s, creating a greater pool  
of the riffle area. Though  
urrence may represent a  
al conditions that existed  
lement in the area. Beaver  
drainage or cause flood-  
due to the water-carrying  
d creek. As noted above,  
ts were common at the  
stream.

ons, especially competi  
community formation and  
t environments Physical  
ly, prevents the continu-  
small group of specialized  
vs rapid colonizers to co-  
stream systems are subject  
nces that fluctuate widely  
of occurrence, including  
lated hydraulic events and  
of materials from the  
am channel. Patch dynam-  
simplifies the rapid coloni-  
that characterizes stream  
inhabit these habitats. In  
ve observed continuing  
ebrate community despite  
more stable riffle-pool

sequence. We specifically attributed the popula-  
tion shifts to (1) environmental fluctuations,  
(2) creation of the new riffle-pool sequences  
with greatly enlarged pools, (3) increased preda-  
tion from fish, and (4) the rapid colonization  
ability of aquatic invertebrates.

The first half of our study appears to measure  
a response to a large-scale event, specifically cat-  
astrophic high flow and the resulting large-scale  
bed substrate movement in 1991. PAYNE &  
MILLER (1991) found that floods and low flows  
have potentially major significance in determin-  
ing the structure and abundance of invertebrate  
communities in streams with erosive sediments.  
The similarity in species richness observed dur-  
ing the first 4 years of the present study indi-  
cates that a recovery was occurring during the  
first 3 years and that the communities had  
recovered by the fourth year. The deviation of  
the rehabilitated stream from the two reference  
streams (which decreased in an almost identical  
manner) during the final year of the study  
could be attributed to an ability of the rehabili-  
tated stream to withstand (and even improve  
during) some regional effect which decreased  
species richness at the reference streams.


In addition to recovery from catastrophic  
flows, rehabilitation construction increased  
pool habitat in Goodwin Creek from 20% to  
70% of the water surface area. The percent of  
sand decreased while the stable substrate for  
invertebrates increased. With this dramatic  
habitat change, the fish community expanded  
rapidly. The fish biomass before rehabilitation  
was 6.8 kg/100 m, after rehabilitation, it was  
10.0 kg/100 m. The mean numbers of fish  
increased from 3,304 fish/100 m to 4,096 fish/  
100 m. Invertebrates attached to stable intro-  
duced rock and natural stream gravel provided  
the major diet component for key fishes. Gut  
content analysis on three common fish species  
found larger following rehabilitation (longear  
sunfish, *Lepomis megalotis*; bluegill, *L. macrochi-  
rus*; and yellow bullhead catfish, *Ameiurus nata-  
lis*) showed that their entire diet was composed  
of substrate oriented invertebrates, except for  
occasional incidental terrestrial invertebrates.  
Chironomidae composed over 70% of the gut  
content in all three species. This explained the

decrease in Chironomidae instead of an  
expected increase. When pools create vulnera-  
bility because prey are conspicuous or lack ref-  
uge, predation is most effective (POWER 1992).  
This basic habitat change also explained some  
of the observed decline in EPT abundance.

Comparisons of pre- and post-rehabilitation  
information showed other notable differences  
between our rehabilitated stream and the two  
comparison streams. Overall abundance of  
invertebrate individuals decreased at the reha-  
bilitated stream, while abundance increased at  
the reference streams. Invertebrate communi-  
ties in the study streams were dominated by  
midges (Diptera: Chironomidae). Identifica-  
tion of subsampled representatives of the Chi-  
ronomidae collected at the incised streams  
showed that pollution-tolerant species of the  
*Polypedilum illinoense* and *Dicrotendipes nervo-  
sus* groups were most abundant. After rehabili-  
tation, the test stream decreased in chironomid  
abundance and had an increase in non-chirono-  
mid individuals. The reference stream increased  
moderately in chironomid abundance and  
greatly increased in the numbers of non-Chi-  
ronomidae, while the incised reference stream  
exhibited a very large increase in Chironomidae  
numbers and only a moderate increase in other  
invertebrates (Table 1). Although the test  
stream had a very high ratio of Chironomidae  
to other individuals before treatment, this ratio  
was similar to that of the reference stream fol-  
lowing rehabilitation.

The % contribution of the dominant taxon  
(PCDT), an accepted measure of a healthy  
community, improved at the rehabilitated  
stream. A high or increased PCDT in a com-  
munity implies that the community is unbal-  
anced or stressed (PIAFKIN et al. 1989); there-  
fore, a large decrease in the PCDT, as seen at  
our rehabilitated stream, implies improvement.  
PCDT at Goodwin Creek decreased by 30%  
following stream modifications. The non-  
incised reference stream exhibited a 23%  
decrease, while the incised reference stream  
PCDT decreased only 6%.

EPT abundance and richness were not mean-  
ingful indicators of success for our study where  
the creation of a pool habitat was sought. Our



Goodwin Creek, Mississippi.

rehabilitated stream exhibited a 50% decrease in EPT individuals following modifications; both the reference streams had increases. The observed decreases in the number of EPT individuals and the number of EPT taxa at Goodwin Creek is most likely explained by changes in the physical habitat (large increases in pool habitat and associated inundation of gravel riffles compounded by beaver activity) and by fish predation which selected against some members of these groups.

The number of collector-filterer group individuals exhibited similar patterns at both the rehabilitated stream and the non-incised reference stream, with the most notable similarity occurring in 1993 when both showed large increases in numbers. Collector-filterers have been shown to be primary colonizers of reclaimed habitat (GORE 1982) and sensitive to sedimentation (LAMBERTI & BERG 1995), and sharp increases in their numbers could be indicative of restored habitat in our stream. Their numbers declined sharply during the second and third year after rehabilitation at Goodwin Creek. This change probably resulted from deeper and slower water characteristics after rehabilitation which decreased the amount of available suspended food particles.

Members of the shredder functional group are sensitive to local and upstream riparian zone effects. Observed increases in the number of shredder functional group individuals during the first 2 years of this project appear attributable to recovery. Following rehabilitation (1993–1995) a similar pattern was observed (though at different scales) at the reference streams. Shredder composition at Goodwin Creek, which had been lower than at the two other creeks, increased greatly following rehabilitation, then exhibited the largest percent contribution of shredder individuals to total individuals of all study streams during 1994 and 1995. We observed that the created riffle-pool sequences allowed better trapping of allochthonous material which favored shredders.

Several indications of stream improvement were observed following rehabilitation of Goodwin Creek: increased taxa richness, much

lower Chironomidae-to-other individuals ratio, a substantially decreased contribution of the dominant taxon, similar changes in the filtering collector communities of the rehabilitated and the non-incised reference stream, and a distinct improvement in the shredder functional group in the rehabilitated stream. The EPT measures of a healthy stream community declined, probably due to physical changes which did not favor these invertebrates, but which promoted a pool habitat inclined toward other invertebrates and certain fish species, especially Centrarchidae (SHIELDS et al. 1995), creating competitive interactions and predation (HARRIS et al. 1995).

### Acknowledgments

Construction work was done by the Corps of Engineers, Vicksburg District, and the USDA Natural Resources Conservation Service. Partial funding for evaluation was received from the Corps of Engineers. Many co-workers from the two construction agencies and the Agricultural Research Service assisted in amassing the data. Their valuable help is appreciated.

### References

- GORE, J. A., 1982. Benthic invertebrate colonization: source distance effects on community composition. *Hydrobiologia* 94, 183–193.
- HARRIS, S. C., MARTIN, T. H., & CUMMINS, K. W., 1995. A model for aquatic invertebrate response to Kissimmee River restoration. *Restoration Ecology* 3(3), 181–194.
- LAMBERTI, G. A. & BERG, M. B., 1995. Invertebrates and other benthic features as indicators of environmental change in Juday Creek, Indiana. *Natural Areas J* 15(3), 249–258.
- PATRICK, R., 1975. Stream communities. In CODY, M. L. & DIAMOND, J. M., (eds) *Ecology and Evolution of Communities*, 445–459. Belknap Press, Cambridge, MA.
- PAYNE, B. S. & MILLER, A. C., 1991. The structure of dominant invertebrate assemblages in a small southeastern stream. *J. Freshwat. Ecol.* 6(3), 257–266.
- PIAFKIN, J. I., BARBOUR, M. T., PORTER, K. D., GROSS, S. K., & HUGHES, R. M., 1989. *Rapid bioassessment protocols for use in streams and rivers: benthic invertebrates and fish*. EPA/444/4-80-001, United States Environmental Protection Agency, Washington, D. C.
- POWER, M. E., 1992. Habitat heterogeneity and the functional significance of fish in river food webs. *Ecology* 73: 1675–1688.
- SHIELDS, F. D., JR., COOPER, C. M. & KNIGHT, S. S., 1992. Rehabilitation of Aquatic Habitats in Unstable Streams. In *Proceedings of the fifth international symposium on river*

*sedimentation* 1093–1100.

many

SHIELDS, F. D., JR., KNIGHT, S. S., 1992. Effects of Channel Incision on Fishes. *Environ. Monit. Assess.* 2: 1–10.

SHIELDS, F. D., JR., KNIGHT, S. S., 1992. Incised Stream Physical Habitat. *Reg. Rivers Res. Manage.* 7: 1–10.

YODZIS, P., 1986. Competition

to-other individuals ratio, increased contribution of the similar changes in the filtering efficiency of the rehabilitated and reference stream, and a distinct shredder functional group in the stream. The EPT measures of the community declined, probably due to changes which did not affect shredders, but which promoted a shift toward other invertebrates, especially Ctenophora (Cotter et al 1995), creating competition and predation (HARRIS et

work done by the Corps of Engineers, and the USDA Natural Resources Service. Partial funding for this work from the Corps of Engineers, and the two construction agencies and the Natural Resources Service assisted in their valuable help is appreciated.

invertebrate colonization source community composition - *Hydrobiologia*

HARRIS, R. H., & CUMMINS, K. W., 1995. Aquatic invertebrate response to Kissimmee River restoration. *Hydrobiologia* 303(3): 181-194

HARRIS, R. H., 1995. Invertebrates and other indicators of environmental change in restored streams. *Natural Areas J* 15(3): 249-258

HARRIS, R. H., 1995. Invertebrate communities - In: CODY, M. L. & HARRIS, R. H. (eds) *Ecology and Evolution of Communities*. Sinauer Press, Cambridge MA

HARRIS, R. H., 1991. The structure of dominance hierarchies in a small southeastern stream. *Hydrobiologia* 6(3): 257-266

HARRIS, R. H., PORTER, K. D., GROSS, S. K., & COOPER, C. M., 1994. *Rapid bioassessment protocols for use in streams: benthic invertebrates and fish* - EPA/600/R-94/010. United States Environmental Protection Agency, Washington DC

HARRIS, R. H., 1994. Heterogeneity and the functional structure of river food webs - *Ecology* 73: 100-110

HARRIS, R. H., COOPER, C. M. & KNIGHT, S. S., 1992. The structure of habitats in unstable streams - In: *Stream habitat: an international symposium on river*

sedimentation 1093-1102 - University of Karlsruhe, Germany

SHIELDS, F. D., JR., KNIGHT, S. S., & COOPER, C. M., 1994. Effects of Channel Incision on Base Flow Stream Habitats and Fishes - *Environ Manage* 18(1): 43-57

SHIELDS, F. D., JR., KNIGHT, S. S., & COOPER, C. M., 1995. Incised Stream Physical Habitat Restoration with Stone Weirs - *Reg Rivers Res Manage* 10: 181-198

YODZIS, P., 1986. Competition, mortality and community

structure - In DIAMOND, J. M. & CASE, T. J. (eds) *Community Ecology* 430-455 - Harper and Row, New York

Authors' address.

CHARLES M. COOPER, SAM TESTA, III, F. D. SHIELDS, JR., U.S. Department of Agriculture, Agricultural Research Service, National Sedimentation Laboratory, P.O. Box 1157, Oxford, MS 38655, USA.